

Instantons in the nonperturbative QCD vacuum

N.O. Agasian* and S.M. Fedorov†

*Institute of Theoretical and Experimental Physics,
117218, Moscow, B. Cheremushkinskaya 25, Russia*

(Dated: 17th September 2002)

The influence of nonperturbative fields on instantons in quantum chromodynamics is studied. Nonperturbative vacuum is described in terms of nonlocal gauge invariant vacuum averages of gluon field strength. Effective action for instanton is derived in bilocal approximation and it is demonstrated that stochastic background gluon fields are responsible for infra-red (IR) stabilization of instantons. Dependence of characteristic instanton size on gluon condensate and correlation length in nonperturbative vacuum is found. Comparison of obtained instanton size distribution with lattice data is made.

PACS numbers: 11.15.Tk, 12.38.Lg

I. INTRODUCTION

Instanton is the first explicit example of nonperturbative (NP) quantum fluctuation of gluon field in QCD. It was introduced in 1975 by Polyakov and coauthors [1]. An important development originated with 't Hooft's classic paper [2], in which he calculated the semiclassical tunneling rate. Instanton gas as a model of QCD vacuum was proposed in pioneer works by Callan, Dashen and Gross [3, 4]. These topologically nontrivial field configurations give a possible explanation of some problems of quantum chromodynamics. Instantons allow to explain anomalous breaking of $U(1)_A$ symmetry and the η' mass [5, 6]. Spontaneous chiral symmetry breaking (SCSB) can be explained with the help of instanton and anti-instanton field configurations in QCD vacuum [7]. An important role of instantons in scalar and pseudoscalar channels was pointed out in paper [8]. Taking into account instantons is of crucial importance for many phenomena of QCD (see [9] and references therein).

At the same time, there is a number of serious problems in instanton physics. The first is the IR divergence of integrals over instanton size ρ at large ρ . This makes it impossible to calculate instantons' contribution to some physical quantities, such as vacuum gluon condensate. Second, "area law" for Wilson loop can not be explained in instanton gas model, hence quasiclassical instanton anti-instanton vacuum lacks confinement which is responsible for hadron spectra. It was recently demonstrated that Casimir scaling which is observed in lattice calculations for the interaction potential between heavy quarks $\bar{Q}Q$ in different representations of $SU(3)$ color group [10] can be violated in instanton gas [11]. This matter is also discussed in [12].

There were many theoretical works aimed to solve the problem of instanton instability in

*Electronic address: agasian@heron.itep.ru

†Electronic address: fedorov@heron.itep.ru

it's size ρ^1 . To some extent they all were based on an attempt to stabilize instanton ensemble due to effects of interaction between pseudoparticles [13]. Today the most popular is the model of "instanton liquid", which was phenomenologically formulated by Shuryak [14, 15]. Using hypothesis that instantons dominate in vacuum averages of local gluodynamical operators, and taking into account dipole-dipole interaction between instanton and antiinstanton, he determined main characteristics of the medium: average distance between pseudoparticles is $\bar{R} \sim 1$ fm and their average size is $\bar{\rho} \sim 1/3$ fm. Thus, a small quantity $\bar{\rho}/\bar{R} \simeq 1/3$ appears and a picture of "granular vacuum" emerges, i.e. vacuum consists of well separated and therefore not very much deformed instantons and anti-instantons. Nevertheless, action of pair interaction $\exp\{S_{int}\} \gg 1$ appears to be important for the structure of instanton ensemble, and therefore one can make a conclusion that instanton ensemble in QCD is not a dilute gas, but an interacting liquid [14].

Quantitatively similar results for parameters of instanton liquid were obtained by Diakonov and Petrov [16]. They have suggested that vacuum in gluodynamics consists of superposition of instantons and antiinstantons, and found that stabilization is related to discovered by them classical repulsion between pseudoparticles. However, further development [17] revealed that instanton ensemble can not be stabilized due to purely classical interaction. Thus, the mechanism for the suppression of large-size instantons in the ensemble of topologically non-trivial fields is still not understood.

A more natural assumption is that there exist other nonperturbative fields apart from instantons in the vacuum, which allow to explain listed above problems. Interaction of small-size instanton $\rho < 0.2$ fm with long-wave gluon fluctuations, described by local vacuum condensate $\langle (gG_{\mu\nu}^a)^2 \rangle$, was studied in [18]. It was demonstrated that those fields lead to even faster growth of instanton density with increasing of it's size ρ . On the other hand, in [19, 20] the study of instantons in stochastic QCD vacuum, parametrized by nonlocal gauge invariant vacuum averages of gluon field $\langle \text{tr } G_{\mu\nu}(x)\Phi(x,y)G_{\sigma\lambda}(y)\Phi(y,x) \rangle$ (here $\Phi(x,y)$ is a parallel transporter) was started. It was shown that standard perturbation theory undergoes a change in NP vacuum [19, 20] and that nonlocal interaction between large-size instanton ($\rho > 1$ fm) and nonperturbative background stops the infrared inflation [20]. Next, it was demonstrated in [21] that nonlocal interaction of instanton with NP vacuum fields stabilizes it on scales of order of correlation length in vacuum condensate.

In the present paper we develop gauge invariant method for calculating effective action of an instanton in the NP vacuum. We show that instanton exists as a stable topologically nontrivial field configuration with characteristic size ρ_c . Value of ρ_c functionally depends on properties of bilocal correlator of NP field $\langle \text{tr } G(x)\Phi(x,y)G(y)\Phi(y,x) \rangle$, i.e. it depends on two quantities: $\langle G^2 \rangle$ – gluon condensate, and "measure of vacuum inhomogeneity" T_g – correlation length in NP vacuum. We don't discuss the problem of instantons distribution in 4D-Euclidean space, i.e. the question about their density N/V . Consideration of this problem in the framework of our approach requires studying interacting instanton-antiinstanton ensemble against the background of nonperturbative fluctuations, described by gauge invariant vacuum averages of gluon field.

This paper is organized as follows. Section II describes general formalism for effective action of an instanton in nonperturbative background. In Section III we study one-loop renormalization of instanton action in the presence of NP fields. It is shown that "perturba-

¹ Callan, Dashen and Gross suggested in their pioneer works [4] to solve some of listed problems with the help of break-down of instantons into meron-antimeron pairs.

tive" part of effective instanton action is "freezed" in the NP background, and approaches constant value when instanton size is large enough $\rho > m_*^{-1} \sim 1\text{GeV}^{-1}$. Section IV is devoted to the study of "direct" interaction between instanton and NP fields. NP fields are parametrized by gauge invariant nonlocal vacuum averages of gluon field (correlators). Using cluster expansion we find effective action for an instanton in bilocal approximation in a gauge invariant way. Results of numerical calculations and discussion of results are given in Section V. Derivation of one-loop renormalization of effective instanton action in nonperturbative vacuum in coordinate space, as well as some mathematical additions are brought out to Appendixes.

II. GENERAL FORMALISM

The influence of NP fluctuations on instanton can be separated into two parts. First, perturbation theory undergoes a change in NP background, which results in the change of standard one-loop renormalization of instanton action. Second, there appears a direct nonlocal interaction of instanton with NP background field.

Standard Euclidean action of gluodynamics has the form

$$S[A] = \frac{1}{2g_0^2} \int d^4x \text{tr} F_{\mu\nu}^2[A] = \frac{1}{4} \int d^4x F_{\mu\nu}^a[A] F_{\mu\nu}^a[A], \quad (1)$$

where $F_{\mu\nu}[A] = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]$ is the strength of gluon field and we use the Hermitian matrix form for gauge fields $A_\mu(x) = g_0 A_\mu^a(x) t^a/2$ and $\text{tr} t^a t^b = \delta^{ab}/2$. We decompose A_μ as

$$A_\mu = A_\mu^{\text{inst}} + B_\mu + a_\mu, \quad (2)$$

where A_μ^{inst} is an instanton-like field configuration with a unit topological charge $Q_T[A^{\text{inst}}] = 1$; a_μ is quantum field (expansion in a_μ reduces to perturbation theory, which in gluodynamics leads to asymptotic freedom); B_μ is nonperturbative background field (with zero topological charge), which can be parametrized by gauge invariant nonlocal vacuum averages of gluon field strength².

In general case effective action for instanton in NP vacuum takes the form

$$Z = e^{-S_{\text{eff}}[A^{\text{inst}}]} = \int [Da_\mu] \left\langle e^{-S[A^{\text{inst}} + B + a]} \right\rangle, \quad (3)$$

where $\langle \dots \rangle$ implies averaging over background field B_μ ,

$$\langle \hat{O}(B) \rangle = \int d\mu(B) \hat{O}(B) \quad (4)$$

and $d\mu(B)$ is the measure of integration over NP fields, explicit form of which is not important for the following consideration.

² In operator product expansion method and in QCD sum rules nonperturbative field is characterized by a set of local gluon condensates $\langle G^2 \rangle$, $\langle G^3 \rangle$, ...

Expanding $S[A]$ up to the second power in a_μ one obtains $Z = \langle Z_1(B)Z_2(B) \rangle$, where

$$Z_1(B) = e^{-S[A^{\text{inst}}]} \int [Da_\mu] \det(\nabla_\mu^2) \exp \left[\frac{1}{g_0^2} \text{tr} \int d^4x \{ -(\nabla_\mu a_\nu)^2 + 2iF_{\mu\nu}[a_\mu, a_\nu] \} \right], \quad (5)$$

$$Z_2(B) = \exp \{ -S[A^{\text{inst}} + B] + S[A^{\text{inst}}] \}. \quad (6)$$

Here we use the notation $\bar{A} \equiv A^{\text{inst}} + B$, $F_{\mu\nu} \equiv F_{\mu\nu}[\bar{A}]$, and $\nabla_\mu a_\nu = \partial_\mu a_\nu - i[\bar{A}_\mu, a_\nu]$ is a covariant derivative. Integration over a and B in (5),(6) corresponds to averaging over fields that are responsible for the physics at different scales. Integration over a_μ takes into account perturbative gluons and describes phenomena at small distances. Averaging over B_μ (formally interaction with gluon condensate) accounts for phenomena at scales of confinement radius. Therefore it is physically clear that averaging factorizes and one obtains

$$Z \rightarrow \langle Z_1(B) \rangle \langle Z_2(B) \rangle. \quad (7)$$

It should be noted that in the limit of infinite number of colors $N_c \rightarrow \infty$ factorization (7) is exact $Z(N_c \rightarrow \infty) = \langle Z_1(B) \rangle \langle Z_2(B) \rangle$. This allows us to write effective action of instanton in NP vacuum as a sum of two terms, "perturbative" and "nonperturbative":

$$S_{\text{eff}}[A^{\text{inst}}] = S_{\text{eff}}^{\text{P}}[A^{\text{inst}}] + S_{\text{eff}}^{\text{NP}}[A^{\text{inst}}], \quad (8)$$

$$S_{\text{eff}}^{\text{P}}[A^{\text{inst}}] = -\ln \langle Z_1(B) \rangle, \quad (9)$$

$$S_{\text{eff}}^{\text{NP}}[A^{\text{inst}}] = -\ln \langle Z_2(B) \rangle = -\ln \langle \exp \{ -S[A^{\text{inst}} + B] + S[A^{\text{inst}}] \} \rangle. \quad (10)$$

III. ONE-LOOP RENORMALIZATION OF INSTANTON ACTION IN NP VACUUM

The general expression for one-instanton field configuration has the well known form

$$A_\mu^{\text{inst}} = 2t^b R^{b\beta} \bar{\eta}_{\mu\nu}^\beta \frac{(x - x_0)_\nu}{(x - x_0)^2} f \left(\frac{(x - x_0)^2}{\rho^2} \right), \quad (11)$$

where

$$R^{b\beta} = 2 \text{tr}(\Omega t^\beta \Omega^\dagger t^b), \quad \Omega \in SU(N_c), \quad (12)$$

$$b = 1, 2, \dots, N_c^2 - 1; \quad \beta = 1, 2, 3.$$

$\bar{\eta}_{\mu\nu}^\alpha$ - 't Hooft symbols, and matrix $R^{b\beta}$ ensures embedding of instanton into $SU(N_c)$ group, $R^{b\beta}$ satisfied relations

$$f^{abc} R^{b\beta} R^{c\gamma} = \varepsilon^{\beta\gamma\delta} R^{a\delta}, \quad R^{b\beta} R^{b\gamma} = \delta^{\beta\gamma}, \quad f^{abc} R^{a\alpha} R^{b\beta} R^{c\gamma} = \varepsilon^{\alpha\beta\gamma}. \quad (13)$$

In singular gauge profile function $f(z)$ satisfies boundary conditions $f(0) = 1$, $f(\infty) = 0$ and the classical solution has the form

$$f(z^2) = \frac{1}{1 + z^2}. \quad (14)$$

The probability to find an instanton is determined by the classical action functional $S_{\text{cl}}[A]$ for the solution (11), (14); that is

$$w \sim \exp \{ -S_{\text{cl}}[A_\mu^{\text{inst}}] \} = \exp \{ -8\pi^2/g_0^2 \}. \quad (15)$$

The preexponential factor was calculated in [2]. The result for one-instanton contribution to the QCD partition function is $Z_1 = \int dn(\rho, x_0, R)$, where dn is the differential instanton density

$$dn(\rho, x_0, R) = [dR]d^4x_0 \frac{d\rho}{\rho^5} d_0(\rho), \quad (16)$$

$$d_0(\rho) = \frac{4.6 \exp\{-1.68 N_c\}}{\pi^2 (N_c - 1)! (N_c - 2)!} \left(\frac{8\pi^2}{g^2(\rho)} \right)^{2N_c} \exp \left\{ -\frac{8\pi^2}{g^2(\rho)} \right\}. \quad (17)$$

In the two-loop approximation in gluodynamics, the coupling constant $g^2(\rho)$ is given by

$$\frac{8\pi^2}{g^2(\rho)} = b \ln \left(\frac{1}{\rho\Lambda} \right) + \frac{b_1}{b} \ln \ln \left(\frac{1}{\rho\Lambda} \right) + O \left(\frac{1}{\ln(1/\rho\Lambda)} \right), \quad b = \frac{11}{3} N_c, \quad b_1 = \frac{17}{3} N_c, \quad (18)$$

$\Lambda \equiv \Lambda_{PV}$ corresponds to the Pauli-Villars regularization scheme, $\Lambda \sim 200$ MeV.

Callan, Dashen and Gross [4] were first to show that in a constant gauge field instanton behaves as a colored four-dimensional dipole. Shifman, Vainshtein and Zakharov [18] generalized this result to the case of interaction between a small-size instanton, $\rho < 0.2$ fm, and nonperturbative long-wave fluctuations that are described by the local vacuum condensate $\langle G^2 \rangle$. As a result, $d_0(\rho)$ is replaced by $d_{\text{eff}}(\rho)$, where

$$d_{\text{eff}}(\rho) \propto (\Lambda\rho)^b \left(1 + \frac{4\pi^4 \langle G^2 \rangle}{(N_c^2 - 1)g^4} \rho^4 + \dots \right). \quad (19)$$

Thus, we arrive at the well-known problem of the infrared inflation of instantons in ρ .

Let us now consider how perturbative renormalization of effective instanton action changes in NP vacuum [20]. We will make use of background field method, developed in [22, 23], and stated in a convenient way by Polyakov [23] when discussing asymptotic freedom in nonabelian gauge theories.

In the absence of the term $\sim F_{\mu\nu}[a_\mu, a_\nu]$, there are four independent components of the field a_μ in expression (5) for Z_1 . Integration with respect to these independent components yields $\left[\det^{-1/2}(\nabla_\mu^2) \right]^4$. Taking into account the ghost determinant in (5), we obtain $\left[\det^{-1/2}(\nabla_\mu^2) \right]^2$. Thus, there are all in all two physical polarizations for the field a_μ instead of four, as it must be. The last term in (5), $\sim F_{\mu\nu}[a_\mu, a_\nu]$ describes the interaction of the external field with the spin of the gluon a_μ .

Hence, there exist two effects. First, there is the motion of charged particles in the external gluomagnetic field $F_{\mu\nu}$. This motion leads to screening similar to orbital Landau diamagnetism. Second, there is the direct interaction of $F_{\mu\nu}$ with the spin of the field a_μ . In the second order in the external field, this interaction yields an antiscreening term, which is analogous to that describing the Pauli paramagnetic effect. With logarithmic accuracy, these two phenomena can be separated. In the absence of the term $\sim F_{\mu\nu}[a_\mu, a_\nu]$ from (5), the expression for $S^P = -\ln Z_1$ then completely corresponds to the effective action of massless scalar electrodynamics with allowance for the isotopic factor. Hence, in the second order in the background field, the "diamagnetic" contribution to S^P can be written as

$$S_{\text{dia}}^P = \int \frac{d^4q}{(2\pi)^4} \Pi_{\mu\nu \text{ dia}}^{ab}(q) \bar{A}_\nu^a(q) \bar{A}_\nu^b(-q), \quad (20)$$

where Π_{dia} is given by the diagrams in Fig. 1.




Figure 1: Diagrams for Π_{dia}

Let us first consider a simpler case of $\bar{A} = A^{\text{inst}}$; that is, we set the nonperturbative field $B_\mu = 0$ and demonstrate how the standard perturbative renormalization of the classical instanton action is reproduced. We then have

$$\Pi_{\mu\nu \text{ dia}}^{ab}(q) = 2f^{acd}f^{bcd}\Pi_{\mu\nu}^{(\text{sc})}(q) = 2N_c\delta^{ab}\Pi_{\mu\nu}^{(\text{sc})}(q) \quad (21)$$

The factor of 2 in (21) stems from two physical polarizations of the field a_μ . The expression for the polarization operator in massless scalar electrodynamics has the form

$$\Pi_{\mu\nu}^{(\text{sc})}(q) = -\frac{1}{4} \int \frac{d^4p}{(2\pi)^4} \frac{(2p+q)_\mu(2p+q)_\nu}{p^2(p+q)^2} + \frac{1}{2}\delta_{\mu\nu} \int \frac{d^4p}{(2\pi)^4} \frac{1}{p^2} = \Pi_1(q^2)(q^2\delta_{\mu\nu} - q_\mu q_\nu). \quad (22)$$

The well-known result for Π_1 in the leading logarithmic approximation is given by

$$\Pi_1(q^2) = \frac{1}{192\pi^2} \ln \frac{\Lambda_0^2}{q^2}, \quad (23)$$

where Λ_0 is the ultraviolet cutoff. Thus, we have

$$\Pi_{\mu\nu \text{ dia}}^{ab}(q) = 2N_c\delta^{ab}(q^2\delta_{\mu\nu} - q_\mu q_\nu)\Pi_1(q^2) \quad (24)$$

Considering that the linear component $F_{\mu\nu}$ satisfies the relation

$$F_{\mu\nu}^a(q)F_{\mu\nu}^b(-q) = 2(q^2\delta_{\mu\nu} - q_\mu q_\nu)\bar{A}_\mu^a(q)\bar{A}_\nu^b(-q) \quad (25)$$

and substituting (24) and (25) into (20), we obtain

$$\begin{aligned} S_{\text{dia}}^{\text{P}} &= \frac{1}{4} \int \frac{d^4q}{(2\pi)^4} \Pi_{\text{dia}}(q^2) F_{\mu\nu}^a(q) F_{\mu\nu}^a(-q), \\ \Pi_{\text{dia}}(q^2) &= 4N_c\Pi_1(q^2) = \frac{N_c}{48\pi^2} \ln \frac{\Lambda_0^2}{q^2} \end{aligned} \quad (26)$$

It should be noted that, in (26), the field strength involves only the linear component $\sim \partial_\mu A_\nu$. However, from gauge invariance and invariance under renormalization-group transformations, it follows that cubic ($\sim A^3$) and fourth-order ($\sim A^4$) terms appear in such a way that they supplement $F_{\mu\nu}^2$ to the standard non-Abelian form.

Following Polyakov [23], we represent the paramagnetic term in the form

$$\begin{aligned}
S_{\text{para}}^{\text{P}} &= -\frac{1}{2g_0^4} \int \frac{d^4q}{(2\pi)^4} F_{\mu\nu}^a(q) F_{\lambda\rho}^b(-q) \times f^{acd} f^{bef} \langle a_\mu^c a_\nu^d(+q) a_\lambda^e a_\rho^f(-q) \rangle \simeq \\
&\simeq -\frac{N_c}{4g_0^4} \int \frac{d^4q}{(2\pi)^4} F_{\mu\nu}^a(q) F_{\mu\nu}^a(-q) \int \frac{d^4q}{(2\pi)^4} \frac{2g_0^4}{p(p+q)^2} = \\
&= \frac{1}{4} \int \frac{d^4q}{(2\pi)^4} \Pi_{\text{para}}(q^2) F_{\mu\nu}^a(q) F_{\mu\nu}^a(-q),
\end{aligned} \tag{27}$$

where

$$\Pi_{\text{para}}(q^2) = -\frac{N_c}{4\pi^2} \ln \frac{\Lambda_0^2}{q^2} \tag{28}$$

Bringing the above results together, we find that one-loop renormalized instanton action is given by

$$S^{\text{P}} = \frac{1}{4} \int \frac{d^4q}{(2\pi)^4} \left(\frac{1}{g_0^2} + \Pi(q^2) \right) F_{\mu\nu}^a(q) F_{\mu\nu}^a(-q) \tag{29}$$

where

$$\begin{aligned}
\Pi(q^2) &= \Pi_{\text{dia}}(q^2) + \Pi_{\text{para}}(q^2) \\
&= \frac{N_c}{\pi^2} \left(\frac{1}{48} - \frac{1}{4} \right) \ln \frac{\Lambda_0^2}{q^2} = -\frac{11}{3} N_c \frac{1}{16\pi^2} \ln \frac{\Lambda_0^2}{q^2}
\end{aligned} \tag{30}$$

As might have been expected, S^{P} involves the effective charge $g^2(q)$ defined at the external field momentum q

$$\frac{1}{g^2(q)} = \frac{1}{g_0^2} - \frac{11}{3} N_c \frac{1}{16\pi^2} \ln \frac{\Lambda_0^2}{q^2} \tag{31}$$

Using the standard renormalization-group arguments, we perform normalization at the momentum

$$\Lambda = \Lambda_0 \exp\left\{-\frac{8\pi^2}{bg^2(\Lambda_0)}\right\} \tag{32}$$

We then have

$$\frac{1}{g^2(q)} = \frac{b}{16\pi^2} \ln \frac{q^2}{\Lambda^2}, \quad b = \frac{1}{3} N_c \tag{33}$$

We consider S^{P} assuming that the background field is an instanton. Because $F_{\mu\nu}^2$ is a gauge-invariant quantity, we use the regular gauge

$$F_{\mu\nu}^a(x) = -4\eta_{\mu\nu}^a \frac{\rho^2}{(x^2 + \rho^2)^2} \tag{34}$$

Considering that the Fourier transform in four-dimensional Euclidean space is given by

$$\tilde{f}(q) = \int d^4x e^{iqx} f(x) = \frac{4\pi^2}{q} \int_0^\infty r^2 dr f(r) J_1(qr), \quad r = |x| \tag{35}$$

we obtain

$$\begin{aligned}
\tilde{F}_{\mu\nu}^a(q) &= \int d^4x e^{iqx} F_{\mu\nu}^a(x) \\
&= -16\pi^2 \eta_{\mu\nu}^a \frac{\rho}{q} \int_0^\infty \frac{r^2 dr}{(r^2 + \rho^2)^2} J_1(qr) = -8\pi^2 \eta_{\mu\nu}^a \rho^2 K_0(\rho q)
\end{aligned} \tag{36}$$

where $K_0(x)$ is a McDonald function. It can easily be shown that the classical action for solution (36) has the standard instanton value

$$\begin{aligned} S_{\text{cl}}[A^{\text{inst}}] &= \frac{1}{4g_0^2} \int \frac{d^4q}{(2\pi)^4} \tilde{F}_{\mu\nu}^a(q) \tilde{F}_{\mu\nu}^a(-q) \\ &= \frac{8\pi^2}{g_0^2} 3\rho^4 \int_0^\infty q^3 dq K_0^2(\rho q) = \frac{8\pi^2}{g_0^2} \\ &\left(\int_0^\infty q^3 dq K_0^2(\rho q) = \frac{1}{3\rho^4} \right) \end{aligned} \quad (37)$$

Substituting (36) into (29), we arrive at

$$S^{\text{P}}[A^{\text{inst}}] = 8\pi^2 3\rho^4 \int_0^\infty q^3 dq K_0^2(\rho q) \frac{1}{g^2(q)}. \quad (38)$$

Let us introduce the dimensionless variable $x = \rho q$ and rewrite (38) in the form

$$S^{\text{P}}[A^{\text{inst}}] = 8\pi^2 3 \int_0^\infty x^3 dx K_0^2(x) \frac{1}{g^2(x/\rho)} \quad (39)$$

where

$$\frac{1}{g^2(x/\rho)} = \frac{b}{16\pi^2} \ln \frac{x^2}{\Lambda^2 \rho^2} = \frac{b}{16\pi^2} \ln \frac{1}{\Lambda^2 \rho^2} + \frac{b}{16\pi^2} \ln x^2 \quad (40)$$

Thus, we finally obtain

$$S^{\text{P}}(\rho) = \frac{8\pi^2}{g^2(\rho)} + \text{const} = b \ln \frac{1}{\Lambda \rho} + \text{const}. \quad (41)$$

We have reproduced the standard expression for $S^{\text{P}}(\rho)$. In the instanton density $d(\rho)$, the constant from (41) appears as a preexponential factor, which is immaterial in our approach. We will now study the modifications that arise in the above equations as the result of taking nonperturbative vacuum fluctuations into account.

It was shown in [24] that, for the polarization operator in the nonperturbative vacuum, the exponential decay of the form $\sim |x - y|^{-4} \exp\{-m_* |x - y|\}$ is obtained in the infrared region instead of the standard power behavior $\sim |x - y|^{-4}$. Here $m_* \simeq 0.75 m_{0^{++}} \sim 1 \text{ GeV}$, and 0^{++} is the lightest glueball (see Appendix A). We will use this fact in analyzing instanton in the nonperturbative vacuum. It can be seen from (9) that averaging is performed over the field B. This means that the polarization operator in (29) is represented by the diagrams in Fig. 2.

Hence, in the expression (29) for the effective action, we must replace the perturbative expression for the polarization operator by the polarization operator $\Pi_{\text{NP}}(q^2)$ in the nonperturbative background field. The latter was calculated in [24] and is given by

$$\Pi_{\text{NP}}(q^2) \simeq \frac{11}{3} N_c \frac{1}{16\pi^2} \ln \frac{q^2 + m_*^2}{\Lambda_0^2} \quad (42)$$

Effective action (29) is then written as

$$S_{\text{eff}}^{\text{P}}[A^{\text{inst}}] = \frac{1}{4} \int \frac{d^4q}{(2\pi)^4} \tilde{F}_{\mu\nu}(q) \tilde{F}_{\mu\nu}^a(-q) \frac{1}{g_{\text{NP}}^2(q^2)}, \quad (43)$$




Figure 2: Diagrams for polarization operator Π_{NP} in the nonperturbative background field

where

$$\frac{1}{g_{\text{NP}}^2(q^2)} = \frac{b}{16\pi^2} \ln \frac{q^2 + m_*^2}{\Lambda^2}. \quad (44)$$

Substituting the Fourier transform (36) of the instanton field into (43) and introducing the variable $x = \rho q$, we obtain

$$\begin{aligned} S_{\text{eff}}^{\text{P}}[A^{\text{inst}}] &= 24\pi^2 \int_0^\infty x^3 dx K_0^2(x) \frac{1}{g_{\text{NP}}^2(x)} \\ \frac{1}{g_{\text{NP}}^2(x)} &= \frac{b}{16\pi^2} \ln \frac{x^2 + m_*^2 \rho^2}{\Lambda^2 \rho^2} \end{aligned} \quad (45)$$

Because $\ln$ is a slowly varying function and because the main contribution to the integral in (45) comes from the region $x \sim 1$ ($K_0(x \gg 1) \simeq \sqrt{\pi/2x} e^{-x}$), we can take $1/g_{\text{NP}}^2(x)$ out of the integral with logarithmic accuracy and set $x = 1$. In this way, we obtain [19, 20]

$$S_{\text{eff}}^{\text{P}}(\rho) = \frac{8\pi^2}{g_{\text{NP}}^2(1)} = \frac{b}{2} \ln \frac{1/\rho^2 + m_*^2}{\Lambda^2} \quad (46)$$

From (46), it follows that the perturbative expression is recovered for small-size instantons ($\rho \ll 1/m_*$) and that, for $\rho > 1/m_*$, we have $S_{\text{eff}}^{\text{P}} \rightarrow \text{const.}$

Derivation of formula (46) in coordinate space is given in Appendix A.

IV. INTERACTION BETWEEN INSTANTON AND NONPERTURBATIVE VACUUM GLUON FIELDS

We consider effect of NP fields on instanton, i.e. we evaluate $\langle Z_2 \rangle$. In this work we make use of the method of vacuum correlators, introduced in works of Dosch and Simonov [25]. NP vacuum of QCD is described in terms of gauge invariant vacuum averages of gluon fields (correlators)

$$\Delta_{\mu_1 \nu_1 \dots \mu_n \nu_n} = \langle \text{tr } G_{\mu_1 \nu_1}(x_1) \Phi(x_1, x_2) G_{\mu_2 \nu_2}(x_2) \dots G_{\mu_n \nu_n}(x_n) \Phi(x_n, x_1) \rangle,$$

where $G_{\mu\nu}$ is gluon field strength, and $\Phi(x, y) = \text{Pexp}(i \int_y^x B_\mu dz_\mu)$ is a parallel transporter, which ensures gauge invariance. In many cases bilocal approximation appears to be sufficient for qualitative description of various physical phenomena in QCD. Moreover, there are indications that corrections due to higher correlators are small and amount to several percent [11, 26].

The most general form of bilocal correlator, which follows from antisymmetry in Lorentz indices, is given by

$$\langle g^2 G_{\mu\nu}^a(x, x_0) G_{\rho\sigma}^b(y, x_0) \rangle = \langle G^2 \rangle \frac{\delta^{ab}}{N_c^2 - 1} \times \\ \times \left\{ \frac{D(z)}{12} (\delta_{\mu\rho} \delta_{\nu\sigma} - \delta_{\mu\sigma} \delta_{\nu\rho}) + \frac{\overline{D}(z)}{6} (n_\mu n_\rho \delta_{\nu\sigma} + n_\nu n_\sigma \delta_{\mu\rho} - n_\mu n_\sigma \delta_{\nu\rho} - n_\nu n_\rho \delta_{\mu\sigma}) \right\}, \quad (47)$$

where

$$G_{\mu\nu}(x, x_0) = \Phi(x_0, x) G_{\mu\nu}(x) \Phi(x, x_0), \quad (48)$$

$n_\mu = z_\mu/|z| = (x - y)_\mu/|x - y|$ is the unit vector, $\langle G^2 \rangle \equiv \langle g^2 G_{\mu\nu}^a G_{\mu\nu}^a \rangle$ and, as it follows from normalization, $D(0) + \overline{D}(0) = 1$.

Connection between functions $D(z)$, $\overline{D}(z)$ and standard functions $D(z)$, $D_1(z)$, which are used in works [25, 26, 27, 28], is discussed in Appendix B.

Functions $D(z)$ and $\overline{D}(z)$ have both perturbative and nonperturbative contributions. We will consider only nonperturbative part, because perturbative one has already been taken into account in $S_{\text{eff}}^{\text{P}}$. Most data about nonperturbative components of these functions come from numerical simulations on lattice. Gluon condensate $\langle G^2 \rangle$ is also determined from lattice calculations, but there exists a widely used estimate for this value based on charmonium spectrum analysis and QCD sum rules $\langle G^2 \rangle \simeq 0.5 \text{ GeV}^4$ [29]. According to the lattice data $D(z)$ and $\overline{D}(z)$ are exponentially decreasing functions $D(z) = A_0 \exp(-z/T_g)$, $\overline{D}(z) = A_1 z \exp(-z/T_g)/T_g$, where T_g is the gluonic correlation length, which was measured on the lattice [27, 30] and estimated analytically [31] to be $T_g \sim 0.2 \text{ fm}$. Besides, according to lattice measurements $A_1 \ll A_0$ ($A_1 \sim A_0/10$). Lattice data from paper of Di Giacomo [28] are presented in table I. $SU(3)$ full stands for chromodynamics with 4 quarks, while $SU(2)$ and $SU(3)$ quenched stand for pure $SU(2)$ and $SU(3)$ gluodynamics, respectively. Bilocal correlator was also measured on the lattice without using cooling method [30], and correlation length was found to be of around $0.1 - 0.2 \text{ fm}$ in quenched approximation.

Gluon condensate $\langle G^2 \rangle = 1.02 \pm 0.1 \text{ GeV}^4$, obtained by Narison [32] basing on sum rules, is in good agreement with lattice calculations [27], $\langle G^2 \rangle = 0.87 \text{ GeV}^4$, for $SU(3)$ QCD.

Table I: Lattice data [28] for correlation length

$T_g, \text{ fm}$	
$SU(2)$ quenched	0.16 ± 0.02
$SU(3)$ quenched	0.22 ± 0.02
$SU(3)$ full	0.34

To evaluate $S_{\text{eff}}^{\text{NP}}$ defined in (10) we use the cluster expansion, which is well known in statistical physics [33]:

$$\langle \exp(x) \rangle = \exp \left(\sum_n \frac{\langle\langle x^n \rangle\rangle}{n!} \right), \quad (49)$$

where $\langle x \rangle = \langle\langle x \rangle\rangle$; $\langle x^2 \rangle = \langle\langle x^2 \rangle\rangle + \langle x \rangle^2$; $\langle x^3 \rangle = \langle\langle x^3 \rangle\rangle + 3\langle x \rangle \langle\langle x^2 \rangle\rangle + \langle x \rangle^3$; ...

One can easily find coefficients in cluster expansion, rewriting it in the following way

$$\begin{aligned}
\langle 1 \rangle &= \langle\langle 1 \rangle\rangle \\
\langle 1 2 \rangle &= \langle\langle 1 2 \rangle\rangle + \langle 1 \rangle \langle 2 \rangle \\
\langle 1 2 3 \rangle &= \langle\langle 1 2 3 \rangle\rangle + \langle 1 \rangle \langle\langle 2 3 \rangle\rangle + \langle 2 \rangle \langle\langle 1 3 \rangle\rangle + \langle 3 \rangle \langle\langle 1 2 \rangle\rangle + \langle 1 \rangle \langle 2 \rangle \langle 3 \rangle \\
&\dots
\end{aligned} \tag{50}$$

Next, we modify expression (10) for $S_{\text{eff}}^{\text{NP}}$ by adding $S[B]$ to the exponent:

$$S_{\text{eff}}^{\text{NP}}[A^{\text{inst}}] = -\ln \langle \exp\{-S[A^{\text{inst}} + B] + S[A^{\text{inst}}] + S[B]\} \rangle. \tag{51}$$

In bilocal approximation this is equivalent to a change of normalization of partition function,³ which is not important for the following consideration.

Nonperturbative part of the effective instanton action $S_{\text{eff}}^{\text{NP}}$ takes the form

$$\begin{aligned}
S_{\text{eff}}^{\text{NP}} &= \langle S[A^{\text{inst}} + B] - S[B] - S[A^{\text{inst}}] \rangle - \\
&\frac{1}{2} \left(\langle (S[A^{\text{inst}} + B] - S[B] - S[A^{\text{inst}}])^2 \rangle - \langle S[A^{\text{inst}} + B] - S[B] - S[A^{\text{inst}}] \rangle^2 \right) + \dots
\end{aligned} \tag{52}$$

We don't go further than $\frac{1}{2} \langle\langle (S[A^{\text{inst}} + B] - S[B] - S[A^{\text{inst}}])^2 \rangle\rangle$ term. Next terms of cluster expansion decrease as $1/g^{2n}$. It is shown in [21] that $g^2(\rho_c) \sim 4$ on the scale of instanton typical size, and therefore $1/g^2$ is a small parameter. Another small parameter is $1/N_c$.

By using Fock-Schwinger gauge $x_\mu A_\mu = x_\mu B_\mu = 0$ (instanton field A_μ^{inst} satisfies it owing to the properties of 't Hooft symbols) and taking into account that

$$\begin{aligned}
S[A^{\text{inst}} + B] - S[B] - S[A^{\text{inst}}] &= \frac{1}{2g^2} \int d^4x \text{tr} \left\{ -([A_\mu^{\text{inst}}, B_\nu] - [A_\nu^{\text{inst}}, B_\mu])^2 \right. \\
&\quad \left. + 2F_{\mu\nu}[A^{\text{inst}}]G_{\mu\nu}[B] - 4i(F_{\mu\nu}[A^{\text{inst}}] + G_{\mu\nu}[B])[A_\mu^{\text{inst}}, B_\nu] \right\}
\end{aligned} \tag{53}$$

one gets

$$S_{\text{eff}}^{\text{NP}} = S_{\text{dia}} + \frac{1}{2} S_{\text{dia}}^2 + S_{\text{para}} + S_1 + S_2, \tag{54}$$

where in bilocal approximation

$$S_{\text{dia}} = -\frac{1}{2g^2} \int d^4x \left\langle \text{tr} \left[([A_\mu^{\text{inst}}, B_\nu] - [A_\nu^{\text{inst}}, B_\mu])^2 \right] \right\rangle, \tag{55}$$

$$S_{\text{para}} = -\frac{1}{2g^4} \int d^4x d^4y \left\langle \text{tr} (F_{\mu\nu}^{\text{inst}}(x) G_{\mu\nu}(x)) \text{tr} (F_{\rho\sigma}^{\text{inst}}(y) G_{\rho\sigma}(y)) \right\rangle, \tag{56}$$

$$S_1 = \frac{2}{g^4} \int d^4x d^4y \left\langle \text{tr} (F_{\mu\nu}^{\text{inst}}[A_\mu^{\text{inst}}, B_\nu])_x \text{tr} (F_{\rho\sigma}^{\text{inst}}[A_\rho^{\text{inst}}, B_\sigma])_y \right\rangle, \tag{57}$$

$$S_2 = \frac{2i}{g^4} \int d^4x d^4y \left\langle \text{tr} (F_{\mu\nu}^{\text{inst}} G_{\mu\nu})_x \text{tr} (F_{\rho\sigma}^{\text{inst}}[A_\rho^{\text{inst}}, B_\sigma])_y \right\rangle. \tag{58}$$

³ Indeed, starting from (10) one finds that in bilocal approximation (54) gets additional terms of the type $\langle G^2 \rangle \int d^4x$, which do not depend on A^{inst} and therefore result only in renormalization of Z_2 .

We use notations S_{dia} (diamagnetic) and S_{para} (paramagnetic) for contributions (55) and (56) into interaction of instanton with background field. Physical motivation for this is discussed in detail in [20].

We also take into account that

$$\langle \text{tr}(G_{\mu\nu}(A_\mu^{\text{inst}} B_\nu - B_\nu A_\mu^{\text{inst}})) \rangle = i \frac{g^3}{2} f^{abc} \langle G_{\mu\nu}^a B_\nu^c \rangle A_\mu^{\text{inst} b} \propto f^{abc} \delta^{ac} = 0. \quad (59)$$

In Fock-Schwinger gauge

$$B_\mu(x) = x_\nu \int_0^1 \alpha d\alpha G_{\nu\mu}(\alpha x), \quad (60)$$

and thus we substitute vacuum averages by correlators in (55)-(58). In general form instanton field has the form

$$A_\mu^{\text{inst}}(x) = \Phi(x, x_0) A_\mu^{\text{sing}}(x - x_0) \Phi(x_0, x). \quad (61)$$

Here $\Phi(x, y) = \text{Pexp} \left(i \int_y^x B_\mu dz_\mu \right)$ is the parallel transporter, and $A_\mu^{\text{sing}}(x - x_0)$ is instanton in fixed (singular) gauge (11),(14). We then obtain for field strength

$$F_{\mu\nu}^{\text{inst}} = \partial_\mu A_\nu^{\text{inst}} - \partial_\nu A_\mu^{\text{inst}} - i[A_\mu^{\text{inst}}, A_\nu^{\text{inst}}] = \Phi(x, x_0) F_{\mu\nu}^{\text{sing}}(x - x_0) \Phi(x_0, x) + \dots \quad (62)$$

Dots denote terms which contain background field B , and therefore lead to higher correlators [20], and we take into account bilocal correlator only.

Parallel transporters along with background field form gauge invariant combinations, which are substituted by correlators after averaging. For instance, one gets for S_{para}

$$\begin{aligned} S_{\text{para}} &= -\frac{1}{2g^4} \int d^4x d^4y \langle \text{tr}(F_{\mu\nu}(x) G_{\mu\nu}(x)) \text{tr}(F_{\rho\sigma}(y) G_{\rho\sigma}(y)) \rangle = \\ &= -\frac{1}{2g^4} \int d^4x d^4y \langle \text{tr}(\Phi(x, x_0) F_{\mu\nu}^{\text{sing}}(x - x_0) \Phi(x_0, x) G_{\mu\nu}(x)) \times \\ &\quad \times \text{tr}(\Phi(y, x_0) F_{\rho\sigma}^{\text{sing}}(y - x_0) \Phi(x_0, y) G_{\rho\sigma}(y)) \rangle = \\ &= -\frac{1}{2g^4} \int d^4x d^4y \langle \text{tr}(F_{\mu\nu}^{\text{sing}}(x - x_0) G_{\mu\nu}(x, x_0)) \text{tr}(F_{\rho\sigma}^{\text{sing}}(y - x_0) G_{\rho\sigma}(y, y_0)) \rangle = \\ &= -\frac{1}{8g^2} \int d^4x d^4y (F_{\mu\nu}^{\text{sing}}(x - x_0))^a (F_{\rho\sigma}^{\text{sing}}(y - x_0))^b \langle g^2 G_{\mu\nu}^a(x, x_0) G_{\rho\sigma}^b(y, x_0) \rangle \end{aligned} \quad (63)$$

Analogous calculations for other contributions to $S_{\text{eff}}^{\text{NP}}$ (54), lead to the following expressions (for the sake of simplicity we omit further index 'sing' for instanton field in singular gauge, $A_\mu^{\text{sing}} \equiv A_\mu$).

$$S_{\text{dia}} = \frac{1}{2} \int d^4x \int_0^1 \alpha d\alpha \int_0^1 \beta d\beta f^{abc} f^{dec} (A_\mu^a A_\mu^d x_\lambda x_\rho \langle g^2 G_{\lambda\nu}^b(\alpha x, x_0) G_{\rho\nu}^e(\beta x, x_0) \rangle - A_\mu^a A_\nu^d x_\lambda x_\rho \langle g^2 G_{\lambda\nu}^b(\alpha x, x_0) G_{\rho\mu}^e(\beta x, x_0) \rangle) \quad (64)$$

$$S_{\text{para}} = -\frac{1}{8g^2} \int d^4x d^4y F_{\mu\nu}^a(x) F_{\rho\sigma}^b(y) \langle g^2 G_{\mu\nu}^a(x, x_0) G_{\rho\sigma}^b(y, x_0) \rangle \quad (65)$$

$$S_1 = -\frac{1}{2} \int d^4x d^4y F_{\mu\nu}^a(x) A_\mu^b(x) F_{\rho\sigma}^d(y) A_\rho^e(y) f^{abc} f^{def} \times \int_0^1 \alpha d\alpha \int_0^1 \beta d\beta x_\xi y_\eta \langle g^2 G_{\xi\nu}^c(\alpha x, x_0) G_{\eta\sigma}^f(\beta y, x_0) \rangle \quad (66)$$

$$S_2 = -\frac{1}{2g} \int d^4x d^4y \int_0^1 \alpha d\alpha F_{\mu\nu}^a(x) F_{\rho\sigma}^b(y) f^{bcd} A_\rho^c(y) y_\xi \langle g^2 G_{\mu\nu}^a(x, x_0) G_{\xi\sigma}^d(\alpha y, x_0) \rangle \quad (67)$$

using representation (47) for gauge invariant condensate $\langle g^2 G_{\mu\nu}^a(x, x_0) G_{\rho\sigma}^b(y, x_0) \rangle$ one gets

$$S_{\text{dia}} = \frac{\langle G^2 \rangle}{12} \frac{N_c}{N_c^2 - 1} \int d^4x \int_0^1 \alpha d\alpha \int_0^1 \beta d\beta x^2 (A_\mu^a(x))^2 [D((\alpha - \beta)x) + 2\overline{D}((\alpha - \beta)x)] \quad (68)$$

$$S_{\text{para}} = -\frac{\langle G^2 \rangle}{48g^2} \frac{1}{N_c^2 - 1} \int d^4x d^4y \left[F_{\mu\nu}^a(x) D(x - y) F_{\mu\nu}^a(y) + 4 \frac{(x - y)_\mu (x - y)_\rho}{(x - y)^2} F_{\mu\nu}^a(x) \overline{D}(x - y) F_{\rho\nu}^a(y) \right] \quad (69)$$

$$S_1 = -\frac{\langle G^2 \rangle}{24} \frac{1}{N_c^2 - 1} \int d^4x d^4y \int_0^1 \alpha d\alpha \int_0^1 \beta d\beta (xy \delta_{\nu\sigma} - x_\sigma y_\nu) f^{abe} f^{cde} \times F_{\mu\nu}^a(x) A_\mu^b(x) F_{\rho\sigma}^c(y) A_\rho^d(y) D(\alpha x - \beta y) + O(\overline{D}) \quad (70)$$

$$S_2 = \frac{\langle G^2 \rangle}{12g} \frac{1}{N_c^2 - 1} \int d^4x d^4y f^{abc} F_{\rho\sigma}^a(y) A_\rho^b(y) F_{\sigma\nu}^c(x) y_\nu \int_0^1 \alpha d\alpha D(x - \alpha y) + O(\overline{D}) \quad (71)$$

We have taken into account, that

$$x_\mu \tilde{B}_\mu(x) \equiv x_\mu (\Phi(x_0, x) B_\mu(x) \Phi(x_0, x)) = \Phi(x_0, x) (x_\mu B_\mu(x)) \Phi(x_0, x) = 0, \quad (72)$$

and thus

$$\tilde{B}_\mu(x) = x_\nu \int \alpha d\alpha \tilde{G}_{\mu\nu}(\alpha x) = x_\nu \int \alpha d\alpha G_{\mu\nu}(\alpha x, x_0) + \dots, \quad (73)$$

where in analogy with (62)

$$\tilde{G}_{\mu\nu}(x) = \partial_\nu \tilde{B}_\mu - \partial_\mu \tilde{B}_\nu - i[\tilde{B}_\mu, \tilde{B}_\nu] = \Phi(x_0, x) G_{\mu\nu}(x) \Phi(x, x_0) + \dots = G_{\mu\nu}(x, x_0) + \dots \quad (74)$$

Thus we have obtained effective action for instanton in NP vacuum in bilocal approximation⁴. It can be seen from numerical calculation that typical instanton size in QCD is $\rho_c \sim 0.25 \div 0.3 \text{ fm}$. For such ρ instanton field in it's center is strong $F_{\mu\nu}^2(x = x_0, \rho_c) = 192/\rho_c^4 \gg \langle G^2 \rangle$, and therefore classical instanton solution is not much deformed in the region $|x| < \rho$, which gives the main contribution to integrals (68)–(71).

The problem of asymptotic behavior of instanton solution far from the center $|x| \gg \rho$ was studied in detail in refs. [16, 34, 35]. Our numerical analysis shows that the value of ρ_c is almost not affected by the asymptotic of classical instanton solution provided that condensate $\langle G^2 \rangle$ and correlation length T_g have reasonable values.

Next we have to define function $D(z)$. We use gaussian form $D(z) = \exp(-\mu^2 z^2)$, $\mu \equiv 1/T_g$. $D(z)$ is a monotonously decreasing function with typical correlation length T_g . Numerical calculations show that explicit form of $D(z)$, as well as taking into account $\bar{D}(z)$ almost do not affect the value of instanton size in NP vacuum ρ_c (see Appendix C).

We use the standard profile function $f = \rho^2/(\rho^2 + x^2)$. Certainly, exact instanton profile (i.e. profile that minimizes action) is different from this one. Nevertheless, knowing the dependence of S_{eff} on ρ we will be able to make a conclusion about instanton behavior (inflation or stabilization) and to determine it's typical size in NP vacuum.

Thus we obtain after integration over space directions

$$S_{\text{dia}} = \frac{2\pi^2 \langle G^2 \rangle}{g^2} \frac{N_c}{\mu^4} \frac{N_c}{N_c^2 - 1} \zeta^4 \int_0^\infty dx \frac{x^3}{(x^2 + \zeta^2)^2} \varphi(x) \quad (75)$$

$$S_{\text{para}} = \frac{-16\pi^4 \langle G^2 \rangle}{g^4} \frac{1}{\mu^4} \frac{1}{N_c^2 - 1} \zeta^4 \int_0^\infty dx dy \frac{x^2 y^2 e^{-(x^2 + y^2)} I_3(2xy)}{(x^2 + \zeta^2)^2 (y^2 + \zeta^2)^2} \quad (76)$$

$$S_1 = \frac{-64\pi^4 \langle G^2 \rangle}{g^4} \frac{1}{\mu^4} \frac{1}{N_c^2 - 1} \zeta^8 \int_0^\infty dx dy \frac{x^2 y^2}{(x^2 + \zeta^2)^3 (y^2 + \zeta^2)^3} \times \\ \times \int_0^1 d\alpha \int_0^1 d\beta e^{-(\alpha^2 x^2 + \beta^2 y^2)} (I_1(2\alpha\beta xy) + I_3(2\alpha\beta xy)) \quad (77)$$

$$S_2 = \frac{-64\pi^4 \langle G^2 \rangle}{g^4} \frac{1}{\mu^4} \frac{1}{N_c^2 - 1} \zeta^6 \int_0^\infty dx dy \frac{x^2 y^2 e^{-x^2}}{(x^2 + \zeta^2)^2 (y^2 + \zeta^2)^3} \int_0^1 d\alpha e^{-\alpha^2 y^2} I_3(2\alpha xy) \quad (78)$$

where $I_n(x) = e^{-\frac{i\pi n}{2}} J_n(ix)$ is Bessel function of imaginary argument (Infeld function), $\zeta \equiv \mu\rho$ and

$$\varphi(x) = \int_0^1 \alpha d\alpha \int_0^1 \beta d\beta e^{-(\alpha - \beta)^2 x^2} = e^{-x^2} \left(\frac{1}{3x^2} - \frac{1}{6x^4} \right) + \frac{1}{6x^4} - \frac{1}{2x^2} + \frac{2}{3x} \Phi(x) \quad (79)$$

where $\Phi(x) = \int_0^x e^{-\xi^2} d\xi$ is error function.

⁴ The tensor structure of instanton was used to derive (68)–(71).

We studied asymptotic behavior of instanton effective action and its dependence on dimensionless parameter $\zeta \equiv \mu\rho$. For a small-size instanton $\rho \ll 1/\mu$ ($\zeta \rightarrow 0$) we found for $S_{\text{eff}}^{\text{NP}}$

$$S_{\text{dia}} \rightarrow -\frac{\pi^2}{2g^2} \frac{N_c}{N_c^2 - 1} \langle G^2 \rangle \rho^4 \ln(\mu\rho), \quad (80)$$

$$S_{\text{para}} + S_1 + S_2 \rightarrow -\frac{2\pi^4}{g^4} \frac{\langle G^2 \rangle}{N_c^2 - 1} \rho^4 + O(\langle G^2 \rangle \rho^4(\mu\rho)). \quad (81)$$

In the opposite limit of large $\rho \gg 1/\mu$ ($\zeta \rightarrow \infty$) one has

$$S_{\text{dia}} \rightarrow \frac{\pi^{7/2}}{6g^2} \frac{N_c}{N_c^2 - 1} \langle G^2 \rangle \frac{\rho^3}{\mu}, \quad (82)$$

$$S_{\text{para}} + S_1 + S_2 \rightarrow \text{const.} \quad (83)$$

Differential instanton density is proportional to $dn/d^4z d\rho \propto \exp(-S_{\text{eff}})$. Thus, “diamagnetic” interaction of instanton with NP fields leads to its infrared stabilization in size ρ .

V. DISCUSSION AND CONCLUSIONS

In this work we have studied instanton behavior in NP vacuum, which is parametrized by gauge invariant vacuum averages of gluon field strength. We have derived effective instanton action in bilocal approximation and have shown that “diamagnetic” term S_{dia} leads to IR stabilization of instanton.

Numerical results for S_{eff} are plotted in Fig. 3. Three lines correspond to $S_{\text{eff}}^{\text{P}}(\rho)$, $S_{\text{eff}}^{\text{NP}}(\rho)$ and $S_{\text{eff}}(\rho) = S_{\text{eff}}^{\text{P}}(\rho) + S_{\text{eff}}^{\text{NP}}(\rho)$ in the case of $SU(3)$ gluodynamics, at $\langle G^2 \rangle = 1.0 \text{ GeV}^4$ and $T_g = 0.3 \text{ fm}$.

In Table II values of ρ_c for different $\langle G^2 \rangle$ and T_g in case of $SU(3)$ gluodynamics are listed. Values of ρ_c in case of QCD with $N_f = 2$ are presented in Table III.

Instanton differential density $dn/d^4z d\rho \sim \exp(-S_{\text{eff}})$ and corresponding lattice data [36] are presented in Fig. 4. All graphs are normalized to the commonly accepted instanton density 1 fm^{-4} . Instanton size $\bar{\rho} \sim 1/3 \text{ fm}$ was first obtained in numerical calculations on the lattice in [37]. It should be mentioned that over the last years the lattice results have not much converged (compare the conferences [38]). Different groups roughly agree on instantons size within a factor of two, e.g. $\bar{\rho} = 0.3, \dots, 0.6 \text{ fm}$ for $SU(3)$ gluodynamics. There is no agreement at all concerning the density N/V . As a tendency, lattice studies give higher density and larger instantons than phenomenologically assumed.

Thus, our results for ρ_c are consistent with phenomenological value $\bar{\rho}$ and lattice data. Furthermore, we can present physical arguments to explain existing deviations with some lattice groups, which give larger than phenomenological values of instanton size. Lattice calculations include cooling procedure, during which some lattice configurations of gluon field are discarded. This procedure can result in a change in gluon condensate $\langle G^2 \rangle$, and thus instanton size distribution is calculated on the lattice at a value of gluon condensate $\langle G^2 \rangle_{\text{cool}}$ which is different from physical value $\langle G^2 \rangle$. It should be noted that uncooled instanton size distribution was studied in the work [39], where authors introduced a new scaling variable (“cooling radius”), which helps to extract information on the uncooled distribution.

We show dependence of ρ_c on $\langle G^2 \rangle$ for several values of T_g in Fig. 5. One can see that increase of $\langle G^2 \rangle$ results in decrease of instanton size, and that effect is a result of nonlocal “diamagnetic” interaction of instanton with NP fields. Fig. 6 shows ρ_c as a function of T_g for several values of $\langle G^2 \rangle$. The smaller T_g , the larger instanton is. It is physically clear that less correlated NP fields ($T_g \rightarrow 0$) have smaller influence on instanton field configuration, which occupies 4D-Euclidean volume with characteristic size ρ ($\rho \gg T_g$). On the other hand, perturbative quantum fluctuations tend to inflate instanton, and therefore ρ_c increases with the decrease of T_g .

We did not go beyond bilocal approximation in this work. As mentioned above, this approximation is good enough not only for qualitative, but also for quantitative description of some phenomena in nonperturbative QCD [26]. In the problem under consideration there are two small parameters. These are $1/g^2(\rho_c) \sim 0.15, \dots, 0.25$ and $1/N_c$, and there powers grow in each term of cluster expansion. Moreover, one can make an estimate of the leading terms in cluster expansion. It follows from our numerical calculations that in the region $\rho \gtrsim \rho_c$ dominating terms are S_{dia} , $\frac{1}{2}S_{\text{dia}}^2$, $\frac{1}{3}S_{\text{dia}}^3$ and so on. Summing them up, one gets $S_{\text{dia}} + \frac{1}{2}S_{\text{dia}}^2 + \frac{1}{3}S_{\text{dia}}^3 + \dots = -\ln(1 - S_{\text{dia}})$. Hence, taking into account next terms of cluster expansion not only conserves IR stabilization, but even leads to some decrease of ρ_c . So, proposed model describes physics of single instanton stabilization in NP vacuum, not only qualitatively, but also quantitatively with rather good accuracy.

Acknowledgements

We are especially grateful to Yu.A. Simonov for stimulating discussions and useful remarks, and we thank E.-M. Ilgenfritz, A.B. Kaidalov, V.A. Novikov and A.V. Yung for discussions of results. The financial support of RFFI grant 00-02-17836 and INTAS grant N 110 is gratefully acknowledged.




Figure 3: Effective action $S_{\text{eff}}^{\text{P}}$ (dotted line), $S_{\text{eff}}^{\text{NP}}$ (dashed line) and $S_{\text{eff}} = S_{\text{eff}}^{\text{P}} + S_{\text{eff}}^{\text{NP}}$ (solid line) ($N_c = 3$, $N_f = 0$, $T_g = 0.3$ fm, $\langle G^2 \rangle = 1.0$ GeV⁴.)




Figure 4: Instanton density $dn/d^4z d\rho$ and lattice data [36] ($N_c = 3$, $N_f = 0$, $T_g = 0.2$ fm, $\langle G^2 \rangle = 0.5$ GeV⁴.)




Figure 5: Instanton size as a function of gluon condensate ($N_c = 3$, $N_f = 0$) at $T_g = 0.1$ fm (dashed line), $T_g = 0.2$ fm (solid line), $T_g = 0.3$ fm (dotted line)




Figure 6: Instanton size as a function of correlation radius ($N_c = 3$, $N_f = 0$) at $\langle G^2 \rangle = 0.5$ GeV⁴ (solid line), $\langle G^2 \rangle = 0.87$ GeV⁴ (dotted line), $\langle G^2 \rangle = 1.0$ GeV⁴ (dashed line)

Table II: ρ_c (fm) for different values of T_g and $\langle G^2 \rangle$, $N_c = 3$, $N_f = 0$

$\langle G^2 \rangle$, GeV ⁴	0.5	1.0	1.5	2.0
T_g , fm				
0.1	0.305	0.256	0.230	0.213
0.15	0.282	0.236	0.213	0.197
0.2	0.268	0.226	0.204	0.189
0.25	0.256	0.219	0.198	0.184
0.3	0.254	0.215	0.194	0.181
0.35	0.250	0.212	0.191	0.178

Table III: ρ_c (fm) for different values of T_g and $\langle G^2 \rangle$, $N_c = 3$, $N_f = 2$

$\langle G^2 \rangle$, GeV ⁴	0.5	1.0	1.5	2.0
T_g , fm				
0.1	0.308	0.258	0.232	0.215
0.15	0.284	0.238	0.215	0.199
0.2	0.270	0.227	0.205	0.191
0.25	0.262	0.221	0.199	0.185
0.3	0.256	0.216	0.195	0.182
0.35	0.252	0.213	0.192	0.179

Appendix A: DERIVATION OF $S_{\text{eff}}^{\text{P}}$ IN COORDINATE SPACE

Let us consider derivation of (46) in coordinate space [19].

We start with expression (5) for partition function, and normalizing it to the case $A^{\text{inst}} = 0$ we obtain effective instanton action in nonperturbative background

$$S_{\text{eff}}[A^{\text{inst}}] = S_{\text{eff}}^{\text{P}}[A^{\text{inst}}] + S_{\text{eff}}^{\text{NP}}[A^{\text{inst}}] \quad (\text{A1})$$

$$S_{\text{eff}}^{\text{P}} = S_{\text{cl}} - \ln \left\langle \frac{\det \nabla^2}{\det \nabla_0^2} \left(\frac{\det D^2}{\det D_0^2} \right)^{1/2} \right\rangle \quad (\text{A2})$$

$$S_{\text{eff}}^{\text{NP}} = - \ln \langle \exp \{ -S[A^{\text{inst}} + B] + S[A^{\text{inst}}] + S[B] \} \rangle \quad (\text{A3})$$

where

$$\begin{aligned} S_{\text{cl}} &= 8\pi^2/g_0^2, \\ \nabla^2 &= (\nabla_\mu[A^{\text{inst}} + B])^2, \\ (D^2)_{\mu\nu} &= -\nabla^2\delta_{\mu\nu} + 2iF_{\mu\nu}[A^{\text{inst}} + B] \end{aligned} \quad (\text{A4})$$

while subscript '0' implies that one should put $A^{\text{inst}} = 0$.

Keeping in (A2) terms quadratic in instanton field and using (61) one has e.g. for the ghost contribution (gluon contribution can be obtained in the same way, final result will contain, as always, two physical gluon polarizations)

$$S_{\text{eff}}^{\text{P}} = \int d^4x d^4y \text{tr} A_\mu^{\text{sing}}(x) \partial_\mu \partial_\nu \tilde{\Pi}(x, y) A_\nu^{\text{sing}}(y) \quad (\text{A5})$$

where explicitly gauge-invariant two-point function $\tilde{\Pi}$ is given by

$$\tilde{\Pi}(x, y) = \langle \Phi_{\text{adj}}(0, x) \nabla_0^{-2}(x, y) \Phi_{\text{adj}}(y, 0) \nabla_0^{-2}(y, x) \rangle, \quad (\text{A6})$$

here $\nabla_0^{-2}(x, y) = \langle x | 1/\nabla_0^2 | y \rangle$ is the ghost propagator. One can distinguish in (A6) a two-gluon correlator with an additional condition that adjoint parallel transporter Φ_{adj} is passing through the instanton position $x_0 = 0$. The mass spectrum of $\tilde{\Pi}$ is bounded from below by the two-gluon correlator without that condition Π_0 , and the latter has as the lowest state 0^{++} glueball with the mass around 1.5 GeV. In the large- N_c approximation $\Pi_0(Q)$ has only poles [24]

$$\Pi_0(q^2) = \sum_{n=1}^{\infty} \frac{C_n}{q^2 + M_n^2}, \quad M_n^2 = 4\pi\sigma_{\text{adj}}n + \text{const} \quad (\text{A7})$$

as in $\bar{q}q$ case. The high-excited states in (A7) yield

$$\begin{aligned} \Pi_0(q^2) &\sim \psi \left(\frac{q^2 + m_0^2}{m_0^2} \right) + \text{const}, \quad \psi(x) = \frac{\Gamma'(x)}{\Gamma(x)} \\ m_0^2 &= 4\pi\sigma_{\text{adj}} \simeq M^2(0^{++}) \end{aligned} \quad (\text{A8})$$

Replacing $\tilde{\Pi}$ by Π_0 in (A5) and performing Fourier transform one has

$$S_{\text{eff}}^{\text{P}} = \frac{1}{4} \int \frac{d^4q}{(2\pi)^4} \left(\frac{1}{g_0^2} + \Pi_0(q^2) \right) \tilde{F}_{\mu\nu}^a(q) \tilde{F}_{\mu\nu}^a(-q), \quad (\text{A9})$$

where we use the notation $\tilde{F}_{\mu\nu}$ for instanton field (62).

One can approximate $\Pi_0(q^2)$ in the whole Euclidean region by [19, 20, 24]

$$\Pi_0(q^2) \simeq \frac{11N_c}{48\pi^2} \ln \frac{q^2 + m_*^2}{\Lambda^2}, \quad (\text{A10})$$

where m_* is connected to m_0 through a relation

$$m_*^2 = m_0^2 e^{-C} \approx 0,56m_0^2, \quad C = -\Psi(1) = 0,577... \quad (\text{A11})$$

Inserting in (A9) $\tilde{F}_{\mu\nu}^a(q) = -8\pi^2 \eta_{\mu\nu}^a \rho^2 K_0(\rho q)$, where $K_0(z)$ is the McDonald function yields

$$S_{\text{eff}}^{\text{P}}(\rho) = 24\pi^2 \int_0^\infty x^3 dx K_0^2(x) \frac{1}{g^2(x)} \simeq \frac{8\pi^2}{g^2(1)} \quad (\text{A12})$$

where we have defined the effective charge

$$\frac{1}{g^2(x)} = \frac{b}{16\pi^2} \ln \frac{x^2 + m_*^2 \rho^2}{\Lambda^2 \rho^2}, \quad b = \frac{11}{3} N_c, \quad (\text{A13})$$

which is obviously the same result as (46).

Appendix B: BILOCAL CORRELATOR

The commonly used parametrization of bilocal correlator [25, 26, 27, 28] is different from (47). It has the form

$$\begin{aligned} \langle \text{tr } G_{\mu\nu}(x) \Phi(x, y) G_{\rho\sigma}(y) \Phi(y, x) \rangle &\sim (\delta_{\mu\rho} \delta_{\nu\sigma} - \delta_{\mu\sigma} \delta_{\nu\rho}) \left[\tilde{D}(z) + \tilde{D}_1(z) \right] \\ &+ (z_\mu z_\rho \delta_{\nu\sigma} + z_\nu z_\sigma \delta_{\mu\rho} - z_\mu z_\sigma \delta_{\nu\rho} - z_\nu z_\rho \delta_{\mu\sigma}) \frac{\partial \tilde{D}_1}{\partial z^2} \end{aligned} \quad (\text{B1})$$

Clearly, one can establish correspondence between our definition and (B1)

$$\begin{aligned} \tilde{D}(z) + \tilde{D}_1(z) &\rightarrow D(z) \\ \frac{1}{2} z^2 \frac{\partial \tilde{D}_1}{\partial z^2} &\rightarrow \overline{D}(z) \end{aligned} \quad (\text{B2})$$

Correlator (B1) was measured on the lattice [27]. Functions $\tilde{D}(z)$ and $\tilde{D}_1(z)$ were approximated as

$$\begin{aligned} \tilde{D}(z) &= A_0 \exp(-|z|/T_g) + \frac{a_0}{z^4} \exp(-|z|/\lambda) \\ \tilde{D}_1(z) &= A_1 \exp(-|z|/T_g) + \frac{a_1}{z^4} \exp(-|z|/\lambda) \end{aligned} \quad (\text{B3})$$

Second term in the right hand side of equations (B3) corresponds to perturbative contribution. Numerical values for T_g , obtained in [27], are presented in the Table I. Moreover, it was shown in these works that ratio A_1/A_0 is a small quantity, $A_1/A_0 \simeq 0.1$. Turning back to our notation (D and $\overline{D}$), we get for nonperturbative part

$$\begin{aligned} D(z) &= (A_0 + A_1) e^{-z/T_g} \\ \overline{D}(z) &= -\frac{1}{4} \frac{z}{T_g} A_1 e^{-z/T_g} \end{aligned} \quad (\text{B4})$$

It is clear that in the region $z \sim T_g$, which gives the main contribution to all quantities, calculated in Section IV, $\overline{D}(z) \ll D(z)$ because $A_1 \ll A_0$.

In our calculations we use gaussian parametrization

$$\begin{aligned} \tilde{D}(z) &= A_0 e^{-z^2/T_g^2} \\ \tilde{D}_1(z) &= A_1 e^{-z^2/T_g^2} \end{aligned} \quad (\text{B5})$$

Thus, we have

$$\begin{aligned} D(z) &= (A_0 + A_1) e^{-z^2/T_g^2} \\ \overline{D}(z) &= -\frac{1}{2} \frac{z^2}{T_g^2} A_1 e^{-z^2/T_g^2} \end{aligned} \quad (\text{B6})$$

and still $\overline{D}(z) \ll D(z)$ in the region $z \sim T_g$.

Appendix C: DEPENDENCE OF S_{dia} ON FUNCTIONS D AND $\bar{D}$

Let us consider numerical changes in effective action due to taking into account function $\bar{D}(z)$ in bilocal correlator (47). As stated above, nonlocal "diamagnetic" interaction leads to instanton stabilization in size. Correspondingly, we will consider changes of S_{dia} due to $\bar{D}$ and to taking exponential parametrization of bilocal correlator. Starting from (68)-(71) and taking into account (B6) one gets for S_{dia} (instead of (75)):

$$S_{\text{dia}} = \frac{2\pi^2}{g^2} \frac{\langle G^2 \rangle}{\mu^4} \frac{N_c}{N_c^2 - 1} \zeta^4 \int_0^\infty dx \frac{x^3}{(x^2 + \zeta^2)^2} \varphi(x),$$

$$\varphi(x) = \int_0^1 \alpha d\alpha \int_0^1 \beta d\beta \left[(A_0 + A_1) e^{-(\alpha-\beta)^2 x^2} - A_1 (\alpha - \beta)^2 x^2 e^{-(\alpha-\beta)^2 x^2} \right], \quad (\text{C1})$$

$$A_0 \simeq 0.9, \quad A_1 \simeq 0.1$$

Fig. 7 shows S_{dia} calculated according to formulas (75)-(79) and (C1). One can see, that the difference is very small, and therefore our approximation, when we neglect $\bar{D}(z)$ as compared to $D(z)$ is justified. In the same figure there also is shown graph for S_{dia} , calculated with $D(z) = e^{-|z|/T_g}$, $\bar{D}(z) = 0$. It is clear that exact form of function $D(z)$ does not significantly influence the result. The only property this function should have is to be monotone decreasing function of z with characteristic correlation length T_g .




Figure 7: S_{dia} , with (dashed line) and without (solid line) taking $\bar{D}(z)$ into account. Dash-dotted line – S_{dia} at $D(z) = \exp\{-|z|/T_g\}$.

-
- [1] A. M. Polyakov, Phys. Lett. **B59**, 79 (1975); A. A. Belavin, A. M. Polyakov, A. S. Shvarts and Y. S. Tyupkin, Phys. Lett. **B59**, 85 (1975).
- [2] G. 't Hooft, Phys. Rev. **D14**, 3432 (1976).
- [3] C. G. Callan, R. F. Dashen and D. J. Gross, Phys. Lett. **B63**, 334 (1976).
- [4] C. G. Callan, R. F. Dashen and D. J. Gross, Phys. Rev. **D17**, 2717 (1978); Phys. Rev. **D19**, 1826 (1979).
- [5] G. 't Hooft, Phys. Rev. Lett. **37**, 8 (1976).
- [6] E. Witten, Nucl. Phys. **B149**, 285 (1979); G. Veneziano, Nucl. Phys. **B159**, 213 (1979).
- [7] D. Diakonov and V. Y. Petrov, Nucl. Phys. **B272**, 457 (1986).
- [8] B. V. Geshkenbein and B. L. Ioffe, Nucl. Phys. **B166**, 340 (1980).
- [9] T. Schafer and E. V. Shuryak, Rev. Mod. Phys. **70**, 323 (1998) [hep-ph/9610451].
- [10] G. S. Bali, Nucl. Phys. Proc. Suppl. **83**, 422 (2000) [hep-lat/9908021]; Phys. Rept. **343**, 1 (2001) [hep-ph/0001312].
- [11] Yu. A. Simonov, JETP Lett. **71**, 127 (2000) [hep-ph/0001244]; V. I. Shevchenko and Yu. A. Simonov, Phys. Rev. Lett. **85**, 1811 (2000) [hep-ph/0001299].
- [12] M. Fukushima, E.-M. Ilgenfritz and H. Toki, Phys. Rev. **D64**, 034503 (2001) [hep-ph/0012358].
- [13] E.-M. Ilgenfritz and M. Muller-Preussker, Nucl. Phys. **B184**, 443 (1981).
- [14] E. V. Shuryak, Nucl. Phys. **B203**, 93 (1982).
- [15] E. V. Shuryak, Phys. Rept. **115**, 151 (1984).
- [16] D. Diakonov and V. Y. Petrov, Nucl. Phys. **B245**, 259 (1984).
- [17] I. I. Balitsky and A. V. Yung, Phys. Lett. **B168**, 113 (1986); A. V. Yung, Nucl. Phys. **B297**, 47 (1988).
- [18] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B163**, 46 (1980).
- [19] N. O. Agasian and Yu. A. Simonov, Mod. Phys. Lett. **A10**, 1755 (1995).
- [20] N. O. Agasian, Phys. Atom. Nucl. **59**, 297 (1996).
- [21] N. O. Agasian and S. M. Fedorov, JHEP **0112**, 019 (2001) [hep-ph/0111208]; N. O. Agasian and S. M. Fedorov, hep-ph/0111305.
- [22] L. F. Abbot, Nucl. Phys. **B184**, 189 (1981).
- [23] A. M. Polyakov, Gauge Fields and Strings. Harwood: Acad. Publ. (1987).
- [24] Yu. A. Simonov, Phys. Atom. Nucl. **58**, 107 (1995) [hep-ph/9311247]; Yu. A. Simonov and J. A. Tjon, Annals Phys. **300**, 54 (2002) [hep-ph/0205165].
- [25] H. G. Dosch, Phys. Lett. **B190**, 177 (1987); H. G. Dosch and Yu. A. Simonov, Phys. Lett. **B205**, 339 (1988); Yu. A. Simonov, Nucl. Phys. **B307**, 512 (1988).
- [26] A. Di Giacomo, H. G. Dosch, V. I. Shevchenko and Yu. A. Simonov, hep-ph/0007223.
- [27] M. Campostrini, A. Di Giacomo and G. Mussardo, Z. Phys. **C25**, 173 (1984); A. Di Giacomo and H. Panagopoulos, Phys. Lett. **B285**, 133 (1992); A. Di Giacomo, E. Meggiolaro and H. Panagopoulos, hep-lat/9603017; M. D'Elia, A. Di Giacomo and E. Meggiolaro, Phys. Lett. **B408**, 315 (1997) [hep-lat/9705032].
- [28] A. Di Giacomo, hep-lat/0012013.
- [29] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147**, 385, 448 (1979).
- [30] G. S. Bali, N. Brambilla and A. Vairo, Phys. Lett. **B421**, 265 (1998) [hep-lat/9709079].
- [31] Yu. A. Simonov, Few Body Syst. **25**, 45 (1998) [hep-ph/9712248]; Yu. A. Simonov, Phys. Atom. Nucl. **61**, 855 (1998).

- [32] S. Narison, Phys. Lett. **B387**, 162 (1996) [hep-ph/9512348].
- [33] Kerzon Huang, Statistical Mechanics, Jon Wiley and Sons, Inc., New York–London (1963); A. Isihara, Statistical Physics, Academic Press, New York–London (1971).
- [34] A. B. Migdal, N. O. Agasian and S. B. Khokhlachev, JETP Lett. **41**, 497 (1985); N. O. Agasian and S. B. Khokhlachev, Sov. J. Nucl. Phys. **55**, 628, 633 (1992); N. O. Agasian, hep-ph/9803252; hep-ph/9904227.
- [35] A. E. Dorokhov, S. V. Esaibegian, A. E. Maximov and S. V. Mikhailov, Eur. Phys. J. **C13**, 331 (2000) [hep-ph/9903450].
- [36] A. Hasenfratz and C. Nieter, Phys. Lett. **B439**, 366 (1998) [hep-lat/9806026].
- [37] M. I. Polikarpov and A. I. Veselov, Nucl. Phys. **B297**, 34 (1988).
- [38] J. W. Negele, Nucl. Phys. Proc. Suppl. **73**, 92 (1999) [hep-lat/9810053]; M. Teper, Nucl. Phys. Proc. Suppl. **83**, 146 (2000) [hep-lat/9909124]; M. Garcia Perez, Nucl. Phys. Proc. Suppl. **94**, 27 (2001) [hep-lat/0011026].
- [39] A. Ringwald and F. Schrempp, Phys. Lett. **B459**, 249 (1999) [hep-lat/9903039].